

Observation of Quantum Jumps in a Single Atom

J. C. Bergquist, Randall G. Hulet, Wayne M. Itano, and D. J. Wineland
Time and Frequency Division, National Bureau of Standards, Boulder, Colorado 80303
 (Received 23 June 1986)

We detect the radiatively driven electric quadrupole transition to the metastable $^2D_{5/2}$ state in a single, laser-cooled Hg II ion by monitoring the abrupt cessation of the fluorescence signal from the laser-excited $^2S_{1/2} \rightarrow ^2P_{1/2}$ first resonance line. When the ion "jumps" back from the metastable D state to the ground S state, the $S \rightarrow P$ resonance fluorescence signal immediately returns. The statistical properties of the quantum jumps are investigated; for example, photon antibunching in the emission from the D state is observed with 100% efficiency.

PACS numbers: 32.80.Pj, 42.50.Dv

Recently, a few laboratories have trapped and radiatively cooled single atoms¹⁻⁴ enabling a number of unique experiments to be performed. One of the experiments now possible is to observe the "quantum jumps" to and from a metastable state in a single atom by monitoring of the resonance fluorescence of a strong transition in which at least one of the states is coupled to the metastable state. When the atomic electron moves to the metastable state, the fluorescence from the strongly driven transition disappears. When the electron drops back into the strongly driven transition, the fluorescence abruptly returns. Thus the strong transition fluorescence is a monitor of the quantum state of the atom. Several years ago, Dehmelt had proposed this optical double-resonance scheme (terming it electron shelving) as an amplification mechanism to detect a weak transition in single-atom spectroscopy.⁵ This technique has been used for several years⁶ in high-resolution spectroscopic studies of samples of many laser-cooled ions, achieving quantum amplifications of 10^6 and higher. In 1981, electron-shelving amplification was used to perform optical-optical double resonance in a single, laser-cooled, trapped ion.² While the signal-to-noise ratio in that experiment was not sufficient to see quantum jumps directly, the fact that the atomic fluorescence would be bistable was noted. More recently, the statistics of quantum switching in a single atom have been theoretically treated in some detail first by Cook and Kimble⁷ and subsequently by several other authors.⁸⁻¹³ In this Letter we report the clear experimental demonstration of quantum jumps in a single laser-cooled $^{198}\text{Hg}^+$ ion stored in a miniature radio-frequency trap.

The basic idea for quantum switching and the associated statistics is illustrated with the three-level system shown in Fig. 1 for the Hg II ion. In Hg II there is a "strong" resonance transition from the $5d^{10}6s^2S_{1/2}$ ground state to the $5d^{10}6p^2P_{1/2}$ state near 194 nm. The lifetime of the $^2P_{1/2}$ state has been measured elsewhere to be 2.3 ± 0.3 ns.¹⁴ Additionally, there is a "weak" electric quadrupole transition from the $^2S_{1/2}$

ground state to the $5d^96s^2D_{5/2}$ state near 281.5 nm. The lifetime of the metastable D state has recently been measured to be about 0.1 s.^{15,16} A laser tuned just below resonance on the highly allowed S - P transition will cool the ion and, in our case, scatter up to 5×10^7 photons/s. By collecting even a small fraction of the scattered photons, we can easily monitor the quantum state of the atom. If only the strong transition is radiatively driven, then for averaging times long compared to the $P_{1/2}$ state lifetime and mean $S \rightarrow P$ excitation time, a steady fluorescence level is expected, corresponding to the atom rapidly cycling between the S and P state. If radiation to drive the weak transition is also admitted, then the atom will occasionally be driven into the metastable state and the fluorescence from the strong transition will abruptly disappear. The cessation of the scattering of many photons on the strong transition for one photon absorbed on the weak transition permits unit detection efficiency of the transition to the metastable state.^{6,17} Some time later, the atom returns spontaneously or is driven out of the D state back to the ground state, which causes a sudden return of the fluorescence on the strong transition. The random on/off "telegraphic" signal provides a direct indication of the quantum state of the ion.

The experimental setup is largely the same as that




FIG. 1. Simplified optical energy-level diagram for Hg II.

used in the two-photon spectroscopic study of the ${}^2S_{1/2}$ - ${}^2D_{5/2}$ transition in HgII reported earlier.¹⁵ In the present experiment, one ion (on some occasions, two ions) was loaded² into the trap and cooled to below 25 mK by 2–5 μ W of sum-frequency-generated radiation near 194 nm (spot size $w_0 \approx 10 \mu\text{m}$) tuned just below the ${}^2S_{1/2}$ - ${}^2P_{1/2}$ first resonance transition. The fluorescence light scattered by the ions was detected at right angles to the 194-nm beam with an overall detection efficiency of about 5×10^{-4} . Peak signal counts exceeded $2 \times 10^4/\text{s}$ against a background counting rate of 200/s. This high counting rate permitted us to monitor the fluorescence at a 10-ms sampling rate with a reasonable signal-to-noise ratio. The fast sampling rate is necessary because of the 100-ms lifetime of the ${}^2D_{5/2}$ state.

Radiation from a frequency-stabilized ring dye laser near¹⁵ 563 nm was doubled to 281.5 nm in order to drive the ${}^2S_{1/2}$ - ${}^2D_{5/2}$ electric-quadrupole-allowed transition directly.¹⁸ The power of the 281.5-nm radiation could be adjusted to as much as 20 μ W. The beam was focused at the center of the trap to a spot size w_0 of approximately 25 μm . A magnetic field of approximately 1 mT (10 G) was applied parallel to the electric field vector of the 281.5-nm radiation and perpendicular to its direction of propagation. The selection rule for the electric-quadrupole-allowed transitions to the various Zeeman states for this configuration is $\Delta m_j = 1$. The frequency of the 281.5-nm radiation was tuned to resonance with the ${}^2S_{1/2} \leftrightarrow {}^2D_{5/2}$ ($m_j = -\frac{1}{2} \leftrightarrow m_j = +\frac{1}{2}$) Zeeman component for the quantum switching results reported here. Also, the resonance signal obtained from the $P_{1/2}$ fluorescence counts during scanning of the 281.5-nm laser over this component revealed a linewidth of less than 8 MHz. If this width is due to Doppler broadening, the ion temperature (in the pseudopotential well) is less than 25 mK and the ion is estimated to be confined to a volume characterized by a linear dimension of less than 0.25 μm .

In order to observe the quantum jumps, we monitor the strong fluorescence at 194 nm while simultaneously admitting the 281.5-nm radiation. A computer strobes and displays the detected 194-nm fluorescence counts accumulated in a counter at 10-ms intervals for running times of 40 s. These data are then stored and the process repeated, but now without the 281.5-nm radiation, or, in some cases, with the 281.5-nm laser detuned from resonance. We repeat the entire sequence numerous times for different power levels of the 281.5-nm light and for different detunings of the 194-nm radiation. An example of 4 s of a sequential set of data with the 281.5-nm radiation (0.3 μ W) first off and then on is shown in Figs. 2(a) and 2(b), respectively.

For the general case of coherent excitation, it is




FIG. 2. Samples of the quantum-jump data. (a) The fluorescence counts detected with no resonance radiation exciting the weak $S_{1/2}$ - $D_{5/2}$ transition. The few jumps observed are likely due to collisions with background Hg atoms and radiative decay to the $D_{3/2}$ state. (b) Both the 194- and 281.5-nm radiation are applied simultaneously. Compared to (a), the interruptions to the fluorescence signal are more frequent. (c) Two ions are trapped and cooled. The interruptions to the fluorescence signal show two levels corresponding to loss of fluorescence from one or both ions. The sampling rate for all the data is 10 ms per point; the length of each sample is 4 s.

necessary to examine the off-diagonal terms of the density matrix in order to include possible coherence effects.^{8,13} However, for the conditions of our experiment, that is, for times longer than the inverse of the excitation and spontaneous emission rates on the strong transition, and when the excitation and emission rates on the strong transition exceed those on the weak transition, the dynamics of the quantum-jump process can be described by effective two-state rate equations for coherent or incoherent excitation.⁸ As a consequence there exist simple probabilities per unit time R_+ and R_- that the electron makes an upward or downward jump on the weak transition. Using the theory of Ref. 8 adapted to arbitrary detuning, and with the additional condition relevant to our experiment that the radiation driving the strong transition is well below saturation, we find that the Einstein A coefficient for the spontaneous decay of the weak transition is related to these probabilities by

$$A({}^2D_{5/2}) \approx R_- - R_+. \quad (1)$$

From data similar to those of Fig. 2(b), we plotted the distribution of off times τ_{off} and the distribution of on times τ_{on} . Theory^{7,8} predicts that the probability density for the time duration of the off (and on) intervals

is given by

$$W_{\text{off}}(\tau_{\text{off}}) = R_- \exp(-R_- \tau_{\text{off}}) \quad (2)$$

for time off, and by

$$W_{\text{on}}(\tau_{\text{on}}) = R_+ \exp(-R_+ \tau_{\text{on}}) \quad (3)$$

for time on. Thus, R_+ and R_- are found from exponential least-squares fits to the data, and from these and Eq. (1) we determine the A coefficient for the metastable $^2D_{2/5}$ state. Unfortunately, the data analysis is complicated by the background events as indicated in Fig. 2(a). Although we have not unambiguously determined their origin, two contributions are collisions with background mercury atoms which temporarily heat the ions² and radiative decay from the $^2P_{1/2}$ state to the $^2D_{3/2}$ state. The lifetime of the $^2D_{3/2}$ state has recently been measured to be about 20 ms.¹⁶ According to theory¹⁹ it decays with nearly equal probability to the lower-lying $^2D_{5/2}$ state and to the ground state. We estimate¹⁷ the probability that a mercury ion in the $^2P_{1/2}$ state will decay to the $^2D_{3/2}$ state as 3×10^{-7} . In the present experiment there is no direct measure of the background pressure at the trap, but estimates based on ion-pump current are consistent with the observed frequency of background events. Estimated recoiling rates are also consistent with the data.

If the background interruptions in the fluorescence signal were due solely to collisions, one would expect their rate of occurrence to be independent of the 194-nm scattering rate. The average duration of the off times after a collision would be a function of the 194-nm intensity and detuning, since these parameters would affect the recoiling rate. If the background interruptions were due solely to radiative decay, one would expect their rate of occurrence to be proportional to the 194-nm scattering rate, but their average duration to be fixed, since this would depend only on the metastable decay rates. The present data indicate that both processes may be present in that both the rate and the average duration of off times vary with scattering rate. Even with these background events, we find the lifetime for spontaneous emission from the $^2D_{5/2}$ level given by Eq. (1) to be 90 ± 30 ms, which is in agreement with the earlier measurements.^{15,16}

Also of interest is the two-time intensity-correlation function⁷⁻¹³ for the 194-nm fluorescence:

$$C(\tau) = \langle I(t)I(t+\tau) \rangle. \quad (4)$$

This expression can be written as⁷

$$C(\tau) = \langle I \rangle^2 + (\langle I^2 \rangle - \langle I \rangle^2) \exp[-(R_+ + R_-)\tau], \quad (5)$$




FIG. 3. Plots of the 194-nm fluorescence intensity correlation function $C(\tau)$ and the 281.5-nm emission correlation function $g_D(\tau)$. Photon antibunching in the 281.5-nm emission is inferred from the observation that $g_D(\tau) \rightarrow 0$ as $\tau \rightarrow 0$.

where

$$\langle I \rangle = I_0 R_- / (R_+ + R_-), \quad (6)$$

$$\langle I^2 \rangle - \langle I \rangle^2 = I_0^2 R_+ R_- / (R_+ + R_-)^2. \quad (7)$$

In Fig. 3 we plot the two-time intensity-correlation function for a typical run. The values of R_+ and R_- obtained from a least-squares fit agree with those derived from the distributions of off and on times.

Other statistical properties can be derived from the quantum switching data. For example, if one neglects the background interruptions to the fluorescence signal, each upward (downward) transition in the fluorescence can be assumed to mark the emission (absorption) of a $^2D_{5/2} \rightarrow ^2S_{1/2}$ photon. Let $g_D(\tau)$ denote the probability that the (assumed) emission of a 281.5-nm photon is followed by the (assumed) emission of another 281.5-nm photon at a time τ later, normalized to 1 at $\tau = \infty$. For our experimental conditions, theory predicts that $g_D(\tau) = 1 - \exp[-(R_+ + R_-)\tau]$, and this is in agreement with the data. In Fig. 3 we plot $g_D(\tau)$ from some of our quantum switching data. The fact that $g_D(\tau) \rightarrow 0$ at $\tau = 0$ implies the existence of photon antibunching^{11,13} in the 281.5-nm radiation from the $^2D_{5/2}$ state. Because of the quantum amplification in the S - P scattering loop the photon antibunching is detected with nearly 100% efficiency.

Finally, in Fig. 2(c) we show quantum switching for the case of two laser-cooled and trapped ions (estimated separation $\approx 2.5 \mu\text{m}$). There are three distinct levels of fluorescence corresponding to (a) a maximum when both ions are in the S - P scattering loop, (b) an intermediate level when one ion is shelved in the D state and only one ion is scattering, and (c) no fluorescence in the rare cases when both ions are shelved in the D state.

In summary, we have demonstrated quantum jumps or switching in a single atom. We have analyzed the statistics and found agreement with earlier published values for the lifetime of the $5d^96s^2D_{5/2}$ state in Hg II. It is interesting to speculate about a single atom in which the upper state on the weak transition is extremely long lived and excited by adiabatic rapid passage.²⁰ In this case the random nature of the excitation is eliminated and one could realize a single-atom switch or flip flop.

The authors are pleased to acknowledge the expert technical assistance of C. Manney during parts of this experiment, and the help of D. J. Larson in the earlier work to obtain single-ion cooling. We are appreciative to H. J. Kimble for helpful discussions of "quantum jumps" and to J. J. Bollinger, R. E. Drullinger, and L. Hollberg for comments on the manuscript. We gratefully acknowledge the support of the U.S. Air Force Office of Scientific Research and the U.S. Office of Naval Research. We note that a Letter by Nagourney, Sandberg, and Dehmelt²¹ describing quantum jumps in a single Ba⁺ atom has appeared subsequent to the submission of our Letter.

¹W. Neuhauser, M. Hohenstatt, P. Toschek, and H. Dehmelt, *Phys. Rev. A* **22**, 1137 (1980).

²D. J. Wineland and W. M. Itano, *Phys. Lett.* **82A**, 75 (1981).

³W. Nagourney, G. Janik, and H. Dehmelt, *Proc. Natl. Acad. Sci. U.S.A.* **80**, 643 (1983).

⁴G. Janik, W. Nagourney, and H. Dehmelt, *J. Opt. Soc. Am.* **B 2**, 1251 (1985).

⁵H. G. Dehmelt, *Bull. Am. Phys. Soc.* **20**, 60 (1975), and

J. Phys. (Paris), Colloq. **42**, C8-299 (1981).

⁶D. J. Wineland, J. C. Bergquist, W. M. Itano, and R. E. Drullinger, *Opt. Lett.* **5**, 245 (1980); D. J. Wineland, W. M. Itano, J. C. Bergquist, J. J. Bollinger, and J. D. Prestage, in *Atomic Physics*, edited by R. S. Van Dyck, Jr., and E. N. Fortson (World Scientific, Singapore, 1985), Vol. 9.

⁷R. J. Cook and H. J. Kimble, *Phys. Rev. Lett.* **54**, 1023 (1985).

⁸H. J. Kimble, R. J. Cook, and A. L. Wells, *Phys. Rev. A* **34**, 3190 (1986).

⁹T. Erber and S. Putterman, *Nature* **318**, 41 (1985).

¹⁰J. Javanainen, *Phys. Rev. A* **33**, 2121 (1986).

¹¹A. Schenzle, R. G. DeVoe, and R. G. Brewer, *Phys. Rev. A* **33**, 2127 (1986).

¹²C. Cohen-Tannoudji and J. Dalibard, *Europhys. Lett.* **1**, 441 (1986).

¹³D. T. Pegg, R. Loudon, and P. L. Knight, *Phys. Rev. A* **33**, 4085 (1986).

¹⁴P. Eriksen and O. Poulsen, *J. Quant. Spectrosc. Radiat. Transfer* **23**, 599 (1980).

¹⁵J. C. Bergquist, D. J. Wineland, W. M. Itano, H. Hemmati, H.-U. Daniel, and G. Leuchs, *Phys. Rev. Lett.* **55**, 1567 (1985).

¹⁶C. E. Johnson, *Bull. Am. Phys. Soc.* **31**, 957 (1986).

¹⁷D. J. Wineland, W. M. Itano, J. C. Bergquist, and F. L. Walls, in *Proceedings of the 35th Annual Symposium on Frequency Control, Philadelphia, 1981* (Electronic Industries Association, Washington, DC, 1981), p. 602.

¹⁸D. J. Wineland, J. C. Bergquist, R. E. Drullinger, H. Hemmati, W. M. Itano, and F. L. Walls, *J. Phys. (Paris), Colloq.* **42**, C8-307 (1981).

¹⁹R. H. Garstang, *J. Res. Natl. Bur. Stand. Sect. A* **68**, 61 (1964).

²⁰See, for example, A. Abragam, *Principles of Nuclear Magnetism* (Clarendon, Oxford, 1961), p. 65.

²¹W. Nagourney, J. Sandberg, and H. Dehmelt, *Phys. Rev. Lett.* **56**, 2727 (1986).